

Filed: 31-Dec-2001

*In the United States Patent and Trademark Office*

**PATENT APPLICATION**

**TITLE:**

**METHOD FOR TREATING GLAUCOMA II B**

**INVENTOR(s):**

Dilip Wagle  
Martin Gall  
Stanley C. Bell  
Edmond J. LaVoie

## METHOD FOR TREATING GLAUCOMA IIB

The present application claims the priority of US Applications 60/296,258, filed 6 June 2001, and 60/259,428, filed 29 December 2000.

The present invention relates to methods for treating glaucoma or improving accommodation (i.e. the process by which the eye adjusts for vision at different distances), and to compounds and compositions for use in such treating. In one aspect, the present invention relates to a method of decreasing the intraocular pressure caused by glaucoma.

Diabetes is the major determinant to the development of visual disability and blindness in parts of the world unencumbered by causes related to malnutrition or infectious diseases. Retinopathy is the leading cause of blindness in diabetics and is a progressive, degenerative disease. Of the many risk factors believed to be associated with diabetic retinopathy, the level of glucose in the plasma has been widely investigated. It is well accepted that a lower incidence of retinopathy is associated with decreased plasma levels of glucose.

Ophthalmologic disorders in diabetes include opacification and glaucoma. As the occurrence of these indications is correlated with the persistent hyperglycemia of the disease. Although the incidence of glaucoma is significant in diabetic populations, glaucoma affects a substantial portion of the general aging population as well.

Primary open angle glaucoma occurs in approximately 4% of diabetics compared to 1.8% of the general population. The reasons for the increase in intraocular pressure that is observed in this disorder are not completely understood. The increase in intraocular pressure that characterizes glaucoma is likely caused by an impairment in the drainage of fluid from the eye at the trabecular meshwork since trabeculectomy restores, at least for a period of time, normal intraocular pressures. The origin of this impairment to fluid movement is currently unknown but may be related to a physical obstruction or restriction to movement of proteins that make up a sieving system in the trabecular meshwork. The trabecular meshwork functions as a sieving system that maintains a restricted flow of intraocular fluid from the eye. The result of excess restriction of this flow is a back pressure that causes increased intraocular pressure.


Replacement of the trabecular meshwork (trabeculectomy) remains an established surgical procedure for improving the filtering of intraocular fluid and for overall reduction of intraocular pressure. This remedy is invasive and of limited effectiveness, since pressure elevation frequently recurs after the procedures.

Current chronic pharmaceutical therapies impose a measure of risk on an already medically compromised patient population. The use of topical B-blockers may affect underlying cardiovascular disease, and carbonic anhydrase inhibitors (e.g. Diamox™) may cause metabolic acidosis. The use of pressure-lowering drugs will be affected by the state of renal disease in compromised elderly and diabetic patients. The drawbacks associated with current pharmaceutical therapies highlight an unmet medical need for a chronic pharmaceutical intervention that is distinct in mechanism of action from current therapies.

New strategies for pharmaceutical intervention in the treatment of glaucoma based upon new mechanisms of action need to be identified. In addition, pharmaceutical agents that decrease the intraocular pressure associated with glaucoma are needed. Also, the methods of improving accommodation provided by the invention allow one to avoid costly and burdensome optical solutions, such as the use of separate reading glasses or glasses with bifocal lenses.

### Summary of the Invention

In one embodiment, the invention relates to a method of treating or ameliorating or preventing glaucoma, decreasing intraocular pressure or improving or ameliorating ocular accommodation in an animal, including a human, comprising administering an intraocular pressure decreasing amount or ocular accommodation improving amount of a compound of the formula I or IA,



wherein the substituent groups are as defined below.

**Detailed Description of the Invention**

In accordance with the present invention a method is provided for the treatment of an animal, preferably a mammal, preferably a human with ophthalmologic disorders including glaucoma and reduced accommodation. Briefly the method of the present invention provides for a method of treatment of mammals with glaucoma or reduced accommodation that can be caused by age or certain age-related diseased states such as diabetes. The method provides for administration of classes of inhibitors of advanced glycation. The invention further provides for methods to monitor the improvement in the ocular condition during the course of the administration of compound.

The agents used in the invention are compounds of formulas I or IA, wherein:

- a. J is oxygen, sulfur, or N-R<sup>d</sup>;
- b. the carbon 2 to nitrogen bond is a double bond except when R<sup>c</sup> is oxo;
- c. the bond between carbons 4 and 5 is a single bond or a double bond (in one embodiment, a single bond);
- d. R<sup>a</sup> and R<sup>b</sup> are
  1. independently selected from hydrogen, acylamino, acyloxyalkyl, alkanoyl, alkanoylalkyl, alkenyl, alkoxy, alkoxy carbonyl, alkoxy carbonylalkyl, alkyl, alkylamino, (C<sub>1</sub>-C<sub>3</sub>)alkylenedioxy, allyl, amino, ω-alkylenesulfonic acid, carbamoyl, carboxy, carboxyalkyl (which alkyl can be substituted with alkyloxyimino), cycloalkyl, dialkylamino, halo, hydroxy, (C<sub>2</sub>-C<sub>6</sub>)hydroxyalkyl, mercapto, nitro, sulfamoyl, sulfonic acid, alkylsulfonfyl, alkylsulfynyl, alkylthio, trifluoromethyl, morpholin-4-yl, thiomorpholin-4-yl, piperidin-1-yl, 4-[C<sub>6</sub> or C<sub>10</sub>] arylpiperidin-1-yl, 4-[C<sub>6</sub> or C<sub>10</sub>] arylpiperazin-1-yl, Ar {wherein, consistent with the rules of aromaticity, Ar is C<sub>6</sub> or C<sub>10</sub> aryl or a 5- or 6-membered heteroaryl ring, wherein the 6-membered heteroaryl ring contains one to three atoms of N, and the 5-membered heteroaryl ring contains from one to three atoms of N or one atom of O or S and zero to two atoms of N, each heteroaryl ring can be fused to a substituted benzene, pyridine, pyrimidine, pyrazine, pyridazine, or (1,2,3)triazine (wherein the ring fusion is at a carbon-carbon double bond of Ar)} or wherein Ar is as above but not heteroaryl fused to pyridine} (in one embodiment, Ar is C<sub>6</sub> or C<sub>10</sub> aryl)}; Ar-alkyl, ArO-, ArSO<sub>2</sub>-, ArSO-, ArS-, ArSO<sub>2</sub>NH-, ArNH, (N-Ar)(N-alkyl)N-, ArC(O)-, ArC(O)NH-, ArNH-C(O)-, and (N-Ar)(N-alkyl)N-C(O)-, or

together R<sup>a</sup> and R<sup>b</sup> comprise methylenedioxy- (in one embodiment, R<sup>a</sup> and R<sup>b</sup> are not acyloxyalkyl, alkenyl, (C<sub>1</sub>-C<sub>3</sub>)alkylenedioxy or allyl, or together do not comprise methylenedioxy); or

2. together with their ring carbons form a C<sub>6</sub>- or C<sub>10</sub>- aryl fused ring; or

5 3. together with their ring carbons form a C<sub>5</sub>-C<sub>7</sub> fused cycloalkyl ring having up to two double bonds including any fused double bond of the containing group, which cycloalkyl ring can be substituted by one or more of the group consisting of alkyl, alkoxy carbonyl, amino, aminocarbonyl, carboxy, fluoro, or oxo (in one embodiment, the ring having no double bonds except any fused double bond of the formula I or IA ring, which cycloalkyl ring can be substituted by one or more of the group consisting of alkyl, amino, aminocarbonyl, carboxy, fluoro, or oxo, where multiple substituents are located on different carbon atoms of the cycloalkyl ring, except in the case of alkyl and fluoro substituents, which can be located on the same or different carbon atoms); or

10 4. together with their ring carbons form a fused 5- or 6-membered heteroaryl ring, wherein the 6-membered heteroaryl ring contains one to three atoms of N, and the 5-membered heteroaryl ring contains from one to three atoms of N or one atom of O or S and zero to two atoms of N; or

15 5. together with their ring carbons form a fused five to eight membered second heterocycle (in one embodiment, a fused five to six membered second heterocycle), wherein the fused heterocycle consists of ring atoms selected from the group consisting of carbon, nitrogen, oxygen, sulfur, and S(O)<sub>n</sub>, wherein n is 1 or 2;

e. R<sup>d</sup> is alkyl, alkenyl, hydrogen, or Ar;

20 f. R<sup>e</sup> is

1. oxo (when  $\Delta^{2,3}$  is not present), or (when  $\Delta^{2,3}$  is present) hydrogen, alkyl, alkylthio, hydrogen, mercapto, amino, amino(C<sub>1</sub>-C<sub>3</sub>)alkyl, or amino(C<sub>6</sub> or C<sub>10</sub>)aryl, or wherein the amino of the last three groups can be substituted with

(a) Ar,

30 (b) Ar-Z-, Ar-alkyl-Z-, Ar-Z-alkyl, Ar-amino-Z-, Ar-aminoalkyl-Z-, or Ar-oxyalkyl-Z-, wherein Z is a carbonyl or -SO<sub>2</sub>-


(c) formyl or alkanoyl (in one embodiment,  $R^c$  is according to one of these optionally substituted three groups; in another, J is S or O (in one embodiment S), and  $R^c$  is hydrogen, oxo, alkyl, amino, amino( $C_1$ - $C_3$ )alkyl or aminophenyl, wherein the amino of the latter three groups can be substituted as outlined here),

2.  $-NHC(O)(CH_2)_n-D-R^f R^f$ , wherein D is oxygen, sulfur or nitrogen, wherein where D is nitrogen n is 0, 1 or 2, but when D is oxygen or sulfur  $n=1$  or 2, and  $R^f$  is present only when D is nitrogen, wherein

(a)  $R^e$  is

(1) Ar,

(2) a group of the formula



wherein E is sulfur, oxygen, or  $N-R^i$ , and  $R^e$ ,  $R^h$  and  $R^i$  are independently the same as  $R^a$ ,  $R^b$  and  $R^d$ , respectively,

(3) a  $C_3$ - $C_8$  cycloalkyl ring having up to one double bond with the proviso that the carbon linking the cycloalkyl ring to D is saturated, which cycloalkyl ring can be substituted by one or more alkyl-, alkoxy-, amino-, aminocarbonyl-, carboxy-, fluoro-, or oxo-substituents;

(4) a 5- or 6-membered heteroaryl ring containing at least one and up to three atoms of N for the 6-membered heteroaryl rings and from one to three atoms of N or one atom of O or S and zero to two atoms of N for the 5-membered heteroaryl rings;

(5) hydrogen, ( $C_2$ - $C_6$ )hydroxyalkyl, alkanoylalkyl, alkyl, alkoxyalkyl, alkenyl, carboxyalkyl (which alkyl can be substituted with alkoxyimino), alkoxyalkyl, a group  $Ar^f$  which is  $C_6$ - or  $C_{10}$ -aryl or a 5- or 6-membered, or 9- or 10-membered

heteroaryl (wherein the heteroatom is one oxygen, one sulfur or one nitrogen) or Ar<sup>b</sup>-alkyl; and

- (b) R<sup>f</sup> is independently hydrogen, (C<sub>2</sub>-C<sub>6</sub>)hydroxyalkyl, alkanoylalkyl, alkyl, alkoxyalkyl, alkenyl, carboxyalkyl (which alkyl can be substituted with alkoxyimino), alkoxyalkyl, Ar<sup>b</sup>, or Ar<sup>b</sup>-alkyl;

wherein aryl, Ar, or Ar<sup>b</sup> can be substituted with, in addition to any substitutions

specifically noted one or more general substituents selected from the group of acylamino, acyloxyalkyl, alkanoyl, alkanoylalkyl, alkenyl, alkoxy, alkoxyalkyl, alkoxyalkylalkyl, alkyl, alkylamino, (C<sub>1</sub>-C<sub>3</sub>)alkylenedioxy, alkylsulfonfyl, alkylsulfynyl, ω-alkylenesulfonic acid, alkylthio, allyl, amino, ArC(O)-, ArC(O)NH-, carboxy, carboxyalkyl, cycloalkyl, dialkylamino, halo, trifluoromethyl, hydroxy, (C<sub>2</sub>-C<sub>6</sub>)hydroxyalkyl, mercapto, nitro, ArO-, Ar-, Ar-alkyl-, sulfamoyl, sulfonic acid, 1-pyrrolidinyl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperazin-1-yl-, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperidin-1-yl, azetidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, piperidin-1-yl (in one embodiment, the substituents are selected from the group of aryl preferred general substituents which are: alkyl, amino, dialkylamino, 1-pyrrolidinyl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperazin-1-yl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperidin-1-yl, azetidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl and piperidin-1-yl); and

heterocycles, except those of Ar and Ar<sup>b</sup>, can be substituted with in addition to any substitutions specifically noted one or more preferred substituents selected from

acylamino, alkanoyl, alkoxy, alkoxyalkyl, alkoxyalkylalkyl, alkyl, (C<sub>1</sub> to C<sub>3</sub>)alkylenedioxy, alkylamino, alkylsulfonfyl, alkylsulfynyl, alkylthio, amino, ArC(O)-, ArO-, Ar-, Ar-alkyl, carboxy, dialkylamino, fluoro, fluoroalkyl, difluoroalkyl, hydroxy, mercapto, oxo, sulfamoyl, trifluoromethyl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperidin-1-yl and 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperazin-1-yl (in one embodiment, the substituents selected from heterocycle preferred general substituents:

acylamino, alkanoyl, alkoxy, alkoxyalkyl, alkoxyalkylalkyl, alkyl, (C<sub>1</sub> to C<sub>3</sub>)alkylenedioxy, alkylamino, alkylsulfonfyl, alkylsulfynyl, alkylthio, amino, ArC(O)-, ArO-, Ar-, Ar-alkyl, carboxy, dialkylamino, fluoro, fluoroalkyl, difluoroalkyl, hydroxy, mercapto, oxo, sulfamoyl, trifluoromethyl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperidin-1-yl and 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperazin-1-yl, wherein multiple

substituents are located on different atoms of the heterocyclic ring, with the proviso that alkyl, alkoxycarbonyl, and fluoro substituents can be substituted on the same carbon atom of the heterocyclic ring);

or a pharmaceutically acceptable salt of said compounds,

- 5           with the proviso that where the compound of formula I is administered to decrease intraocular pressure at least one compound of formula I administered in effective amount is not a thiazole substituted on a ring carbon sulfonamide (the amide of which can be substituted) that has carbonic anhydrase inhibiting activity.

- 10           Primary open angle glaucoma is characterized by an increase in intraocular pressure. The condition of open angle glaucoma is characterized by an increase in the pressure within a person's eye or eyes, called the intraocular pressure. The normal pressure is about 15 mmHg. Elevated pressures of 20-30 mm Hg create a strong risk of damage to the optic nerve and blindness.

- 15           Glucose reacts with proteins by a non-enzymatic, post-translational modification process called non-enzymatic glycosylation. The resulting sugar-derived adduct, the advanced glycosylation end product (AGE), matures to a molecular species that is reactive, and can readily bond to amino groups on adjacent proteins, resulting in the formation of AGE cross-links between proteins.

- 20           It has now been found that certain compounds that inhibit the formation of such sugar-derived adducts, or in some cases are believed to deactivate such adducts or break resulting crosslinks, can reduce intraocular pressure or ameliorate a trend towards elevated pressure.

- 25           Structural matrix proteins isolated from tissues of diabetics and aged individuals are more highly crosslinked than those from nondiabetics or younger individuals and are more resistant to both enzymatic and chemical hydrolysis *in vitro*. It is this cross-linked state of proteins that is believed to cause stiffness of tissues. The cleavage of AGE cross-links between proteins can provide a mechanism-based therapy for restoration of normal tissue function. An agent that cleaves AGE cross-links between proteins or  
30           inhibits their formation can restore more normal sieving function and movement to the trabecular meshwork.

In accordance with the present invention, methods for administering pharmaceutical compositions containing compounds have been developed for the



treating glaucoma, intraocular pressure associated with glaucoma, and reduced accommodation. These agents are either substituted thiazole, oxazole, or imidazole agents as shown in the Summary section above.

- As is noted in the formula for I and IA, the invention includes aromatic thiazole, oxazole, and imidazole analogs, as well as non aromatic analogs thereof such as thiazoline, thiazolidine, oxazoline, oxazolidine, imidazoline, and imidazolidine analogs.

The alkyl, and alkenyl groups referred to above include both C1 to C6 linear and branched alkyl and alkenyl groups, unless otherwise noted. Alkoxy groups include linear or branched C1 to C6 alkoxy groups, unless otherwise noted.

- 10 "Ar" (consistent with the rules governing aromaticity) refers to a C<sub>6</sub> or C<sub>10</sub> aryl, or a 5 or 6 membered heteroaryl ring. The heteroaryl ring contains at least one and up to three atoms of N for the 6 membered heteroaryl ring. The 5 membered heteroaryl ring contains; (1) from one to three atoms of N, or (2) one atom of O or S and zero to two atoms of N. Nonlimiting examples of heteroaryl groups include: pyrrolyl, furanyl, 15 thienyl, pyridyl, oxazolyl, pyrazolyl, pyrimidinyl, and pyridazinyl.

"Ar" can be fused to either a benzene, pyridine, pyrimidine, pyridazine, or (1,2,3) triazine ring.

As used herein, C<sub>6</sub> or C<sub>10</sub> aryl groups and heteroaryl containing five or six, or nine to ten ring members are monocyclic or bicyclic.

- 20 In certain embodiments of the invention, the thiazoles, imidazoles, and oxazoles of the invention contain R<sup>a</sup> and R<sup>b</sup> substitutions that together with their ring carbons (the C4-C5 carbons of the thiazoles, imidazoles, and oxazoles) form a five to eight membered fused heterocycle (i.e. a bicyclic heterocycle is formed). In these embodiments the fused heterocycle is preferably not aromatic. Particular compounds within these embodiments 25 contain sulfur atoms in the fused heterocycle (the ring fused to the thiazoles, imidazoles, and oxazoles). These sulfur atoms in these particular compounds can exist in various oxidation states, as S(O)<sub>n</sub>, where n is 0, 1, or 2.

- In certain embodiments of the invention, thiazoles, imidazoles, and oxazoles of the invention contain R<sup>a</sup> and R<sup>b</sup> substitutions that together with their ring carbons (the 30 C4-C5 carbons of the thiazoles, imidazoles, and oxazoles) form a C5 to C7 cycloalkyl ring having up to double bonds including the C4-C5 double bond. In other embodiments a cycloalkyl ring is present when R<sup>c</sup> is a C3 to C8 cycloalkyl ring. The cycloalkyl groups can be substituted by one or more of the group consisting of alkyl-,

- alkoxycarbonyl-, amino-, aminocarbonyl-, carboxy-, fluoro-, or oxo- substituents. One of ordinary skill in the art will recognize that where cycloalkyl groups contain double bonds, the  $sp^2$  hybridized carbon atoms can contain only one substituent (which cannot be amino- or oxo-).  $Sp^3$  hybridized carbon atoms in the cycloalkyl ring can be geminally substituted with the exception that (1) two amino groups and (2) one amino and one fluoro group can not be substituted on the same  $sp^3$  hybridized carbon atom.

- In certain embodiments of the invention, the thiazoles, imidazoles, and oxazoles of the invention contain  $R^a$  and  $R^b$  substitutions that together with their ring carbons (the C4-C5 carbons of the thiazoles, imidazoles, and oxazoles) form a five or six membered heteroaryl ring (i.e., a bicyclic aromatic heterocycle is formed). A preferred bicyclic aromatic heterocycle of the invention is a purine analog [J is  $N-R^d$  and  $R^a$  and  $R^b$  together with their ring carbons (the C4 and C5 of the imidazole ring) form a pyrimidine ring].

- Aryl, Ar, or  $Ar^\phi$  can be substituted with, in addition to any substitutions specifically noted one or more substituents selected from the group of acylamino, acyloxyalkyl, alkanoyl, alkanoylalkyl, alkenyl, alkoxy, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkylamino, (C1-C3)alkylenedioxy, alkylsulfonyl [alkylSO<sub>2</sub>-], alkylsulfinyl [alkylSO-],  $\omega$ -alkylenesulfonic acid [-alkylSO<sub>3</sub>H where n = 1 to 6], alkylthio, allyl, amino, ArC(O)-, ArC(O)NH-, carboxy, carboxyalkyl, cycloalkyl, dialkylamino, halo, trifluoromethyl, hydroxy, (C2-C6)hydroxyalkyl, mercapto, nitro, ArO-, Ar-, Ar-alkyl-, sulfamoyl, sulfonic acid, morpholin-4-yl, thiomorpholin-4-yl, piperidin-1-yl, 1-pyrrolidinyl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperidin-1-yl and 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperazin-1-yl.

- Heterocycles, except those of Ar and  $Ar^\phi$ , can be substituted with in addition to any substitutions specifically noted one or more substituents selected from acylamino, alkanoyl, alkoxy, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, (C1 to C3)alkylenedioxy, alkylamino, alkylsulfonyl, alkylsulfinyl, alkylthio, amino, ArC(O)-, ArO-, Ar-, Ar-alkyl, carboxy, dialkylamino, fluoro, fluoroalkyl, difluoroalkyl, hydroxy, mercapto, oxo, sulfamoyl, trifluoromethyl, 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperidin-1-yl and 4-[C<sub>6</sub> or C<sub>10</sub>]aryl piperazin-1-yl, wherein multiple substituents are located on different atoms of the heterocyclic ring, with the proviso that alkyl, alkoxycarbonyl, and fluoro substituents

can be substituted on the same carbon atom of the heterocyclic ring. Heterocycles can be substituted with one or more substituents.

The halo atoms can be fluoro, chloro, bromo or iodo. Chloro and fluoro are preferred for aryl substitutions.

5 In some embodiments of this invention, the compounds of formula (I) can form biologically and pharmaceutically acceptable salts. Useful salt forms include the halides (particularly bromides and chlorides), tosylates, methanesulfonates, brosylates, fumarates, maleates, succinates, acetates, mesitylenesulfonates, and the like. Other related salts can be formed using similarly non-toxic, biologically or pharmaceutically  
10 acceptable anions.

Representative, non-limiting examples of compounds of the present invention are:

Thiazole

4,5-Dimethylthiazole

4-Methylthiazole

5-Methylthiazole

4-Methyl-5-(2-hydroxyethyl)thiazole

4-Methyl-5-vinylthiazole

Benzothiazole

2-Aminobenzothiazole

2-Amino-4-chlorobenzothiazole

2-Amino-6-chlorobenzothiazole

2,6-Diamino-benzothiazole

2-Aminothiazole

2,4,5-Trimethylthiazole

2-Amino-5-methylthiazole

2-Amino-4-methylthiazole

2-Acetylthiazole

2-Ethyl-4-methylthiazole

Ethyl 2-(Formylamino)-4-thiazoleacetate

2-(Formylamino)-alpha-(methoxyimino)-4-thiazoleacetic acid

2-Amino-4-phenylthiazole hydrochloride monohydrate

2-Isobutylthiazole

2-Methyl-2-thiazoline

2-Methyl-2-oxazoline

2-Oxazolidone


2-Amino-4-thiazoleacetic acid

5 1-(Thiazolyl)-3-phenyl-urea

1-(Thiazolidinyl)-3-(4-fluorophenyl)-urea

(4-fluorophenyl)thiazolin-2-ylamine

2-(4,6-dimethylpyrimidin-2-ylthio)-N-(1,3-thiazol-2'yl)acetamide, also known as  
 N-(Thiazolyl)-2-(4,6-dimethyl-pyrimidin-2-yl-thio)-acetamide



2-(4-propylphenoxy)-N-(thiazol-2-yl)acetamide

2-furyl-N-[4-(6-methylbenzothiazol-2-yl)phenyl]carboxamide

2-(3,5-Dimethylphenoxy)-N-thiazol-2-yl)acetamide

5,5-Dimethyl-2-(2-naphthylamino)-4,5,6-trihydrobenzothiazol-7-one

15 Imidazole

1-Methylimidazole

1-Ethylimidazole

1-Butylimidazole

1-Vinylimidazole

20 1-Allylimidazole

1-(Trimethylsilyl) imidazole

1-(3-Aminopropyl) imidazole

1-Benzyl imidazole

1-Phenyl imidazole

25 1,5-Dicyclohexyl imidazole

1-(p-Toluenesulfonyl) imidazole

N-Benzoyl-imidazole

4-Methyl-imidazole

4'-(Imidazol-1-yl)-acetophenone

4-(Imidazol-1-yl)-phenol

1-(4-Methoxyphenyl)-1H-imidazole

Methyl-4-(1H-imidazol-1-yl)benzoate

5 1-Methylbenzimidazole

as well as other biologically and pharmaceutically acceptable salts thereof.

Where one or more compounds of formula I are administered to decrease intraocular pressure, at least one compound of formula I administered in effective amount is not a thiazole substituted on a ring carbon sulfonamide (the amide of which can be substituted) that has carbonic anhydrase inhibiting activity. Of course, the composition can include an effective amount of a first agent, as well as a carbonic anhydrase-inhibiting effective amount of another agent, including one of those distinguished above.

- 15 Compounds of the formula I can be conveniently prepared by chemical syntheses well-known in the art. Certain of the compounds are well-known and readily available from chemical supply houses or can be prepared by synthetic methods specifically published therefor. For instance, 4,5-Dimethylthiazole, 4-Methylthiazole, 5-Methylthiazole, 4-Methyl-5-thiazoleethanol, 4-Methyl-5-vinylthiazole, Benzothiazole, 2-Aminobenzothiazole, 2-Amino-4-chlorobenzothiazole, 2-Amino-6-chlorobenzothiazole, 2-Aminothiazole, 2,4,5-Trimethylthiazole, 2-Amino-5-methylthiazole, 2-Amino-4-methylthiazole, 2-Acetylthiazole, 2-Ethyl-4-methylthiazole, Ethyl 2-(Formylamino)-4-thiazoleacetate, 2-(Formylamino)-alpha-(methoxyimino)-4-thiazoleacetic acid, 2-amino-4-phenylthiazole hydrochloride monohydrate, 2-
- 20 Isobutylthiazole, 2-Methyl-2-thiazoline, 2-Methyl-2-oxazoline, 2-Oxazolidone, Thiomorpholine, 2-Amino-4-thiazoleacetic acid, Imidazole, 1-Methylimidazole, 1-Butylimidazole, 1-Vinylimidazole, 1-Allylimidazole, 1-(Trimethylsilyl)imidazole, 1-(3-Aminopropyl)imidazole, 1-Benzylimidazole, 1-Phenylimidazole, 1,5-Dicyclohexylimidazole, 1-(p-Toluenesulfonyl)imidazole, N-Benzoyl-imidazole, 4-Methyl-imidazole,
- 30 4'-(Imidazol-1-yl)-acetophenone, 4-(Imidazol-1-yl)-phenol, 1-(4-Methoxyphenyl)-1H-imidazol and 1-Methylbenzimidazole can be obtained from Sigma (St. Louis, MO), Aldrich (Milwaukee, WI) or Fluka (Milwaukee, WI) (all divisions of Sigma-Aldrich Co.).

1-ethylimidazole can be obtained from TCI America (Portland, OR). N-(Thiazolidinyl)-4-fluoroaniline, N-(Thiazolyl)-2-(4,6-dimethyl-pyrimidin-2-yl-thio)-acetamide, N-(Thiazolyl)-2-(4-propylphenoxy)-acetamide, 2-[4-(N-Furoyl)aminophenyl]-6-methylbenzothiazole,


- 5 N-(Thiazolyl)-2-(3,5-dimethylphenoxy)-acetamide and 2-[(N-(2-Napthalenyl)amino)-[2,3:5,4]-(5,5-dimethyl-cyclohexanonyl)]thiazole can be purchased from MDD, Inc. (Acton, Ontario), a successor to Ortech Corporation.

In one synthetic process to prepare compounds of the general formula I, a thiazole is reacted with an alkyl or acyl halide in the presence of base such as triethylamine, to produce the corresponding alkyl or acyl derivative at the 2 carbon. *See*,  
 10 Medici et al., *J. Org. Chem.* 49: 590-596, 1984. In some cases, a chromatographic step is applied to separate additions at the 2 and 3 positions of the thiazole ring.

In another synthesis of compounds of the formula I wherein R<sup>c</sup> is amino, nitro-containing analogs of compounds of the invention or precursors thereof are  
 15 catalytically hydrogenated to the corresponding amino compounds.

2-Amino thiazole compounds wherein (R<sup>c</sup> is amino) can also be synthesized by reacting thiourea (which can be substituted on at least one amine) with an alpha-halo ketone using the method described in *Vogel's Textbook of Practical Organic Chemistry*,  
 5<sup>th</sup> Edition, John Wiley & Sons, New York, p. 1153 Such a reaction is exemplified by a  
 20 synthesis of 2-amino-4-phenylthiazole:

#### Scheme 1




- N-aryl imidazoles can be prepared using appropriate aromatic nucleophilic  
 25 displacement reactions. For example, fluoro phenyl compounds such as 4-fluorobenzoic acid methyl ester can be used to substitute on the N<sup>1</sup> nitrogen of imidazole to make methyl-4-(1H-imidazol-1-yl)benzoate. *See*, Morgan et al., *J. Med. Chem.* 33: 1091-1097, 1990.


Amino functions of 2-aminoimidazoles or 2-aminothiazoles can be acylated by dehydration or other methods known in the art.

Substituted oxazoles can be prepared by methods known in the art. For instance, 2-unsubstituted oxazoles can be formed by condensation of formamide with either  $\alpha$ -hydroxy or  $\alpha$ -haloketones intermediates (H. Brederick, R. Gommper, H. G. v. Shuh and G. Theilig, in *Newer Methods of Preparative Organic Chemistry*, Vol. III, ed. W. Foerst, Academic press, New York, 1964, p. 241). The intermediates can cyclize under acid conditions to form the oxazole ring (**Scheme 2**). In addition, 2,4-disubstituted oxazoles can be prepared from  $\alpha$ -haloketones and amides at higher temperatures using the same method.



**Scheme 2**



Oxazoles can be prepared by cyclization reactions of isonitriles (van Leusen, A. M. *Lect. Heterocycl. Chem.* **1980**, 5, S111; Walborsky, H. M.; Periasamy, M. P. in *The Chemistry of Functional Groups, suppl. C*, Patai, S.; Rappoport, Z., Eds; Wiley-Interscience, 1983, p. 835; Hoppe, D. *Angew. Chem. Int. Edn. Engl.*, **1974**, 13, 789; Schollkopf, U. *Angew. Chem. Int. Ed. Engl.*, **1977**, 16, 339). For example, as shown below in **Scheme 3**, the tosylmethyl isocyanide can be deprotonated by a base and reacted with a suitable electrophile (e.g. an aldehyde). The intermediate can cyclize and aromatize to provide the desired oxazole analog. Other methods for preparing oxazoles include 1,5-dipolar cyclization of acylated nitrile ylides (Taylor E. C.; Turchi, I. J. *Chem. Rev.*, **1979**, 79, 181; Huisgen, R. *Angew. Chem. Int. Edn. Engl.* **1980**, 19, 947)

**Scheme 3**


2-Amino-substituted oxazoles (i.e.  $\text{R}^c=\text{NH}_2$ ) can be prepared by two general methods. Urea can be condensed with  $\alpha$ -bromo ketones to yield 2-aminooxazoles that can be substituted at the 4 and 5 positions (**Scheme 4**). Alternatively, another route to 2-aminooxazoles from acyclic precursors is the base catalyzed reaction of cyanamide with  $\alpha$ -hydroxy ketones (**Scheme 5**). 2-Aminooxazoles of the invention can also be prepared from the nucleophilic displacement of amines with 2-chlorooxazole, for example, for compounds of the invention wherein  $\text{R}^c$  is  $\text{ArNH-}$  (Gompper, R.; Effenberger, F. *Chem. Ber.*, 1959, 92, 1928).

**Scheme 4****Scheme 5**

Compounds of the invention, wherein  $\text{R}^c$  is arylcarbonyl can be synthesized by acylation of the amino moiety of 2-aminooxazoles with, for example, with anhydrides to yield 2-acylamino oxazoles. In addition, 2-aminooxazoles can be acylated, for instance, with chloroacetic anhydride to yield an  $\alpha$ -chloro carboxamide. The  $\alpha$ -chloro carboxamide can serve as a suitable alkylating reagent that can be treated with, for example, phenols, arylamines and alkylamines to prepare compounds of the invention. An oxazole of the invention wherein  $\text{R}^c$  is Ar-oxycarbonylamino is shown in **Scheme 6**.




Scheme 6




Oxazoles of the invention with R<sub>c</sub> is aminocarbonylamino (ureido) or aminothiocarbonylamino (thioureido) can be prepared from 2-aminooxazoles (**Scheme 7**). 2-Aminooxazoles can be treated with isocyanates and isothiocyanates to yield 2-ureido and 2-thioureido oxazoles, respectively (Crank, G.; Foulis, J. *J. Med. Chem.*, **1971**, *14*, 1075; Crank, G. Neville, M.; Ryden, R. *J. Med. Chem.*, **1974**, *16*, 1402).

Scheme 7




2-Aminooxazoles can be hydrogenated using palladium catalysts to yield 2-aminooxazolines (**Scheme 8**) (Tanaka, C.; Kuriyama, S. *Yakugaku Zasshi* **1979**, *99*, 78).

Scheme 8



Benzoxazole intermediates substituted at the 2 position can be prepared from 2-aminophenols by acylation with, for example, with an acid chloride and cyclization (**Scheme 9**).

Scheme 9



To treat glaucoma or reduced accommodation and associated symptoms, an effective amount of a pharmaceutical compound will be recognized by clinicians but includes an amount effective to treat, reduce, ameliorate, eliminate or prevent one or more symptoms of the disease sought to be treated or the condition sought to be avoided or treated, or to otherwise produce a clinically recognizable change in the pathology of the disease or condition.

In treating glaucoma, agents of the inventions can be administered concurrently or in a combined formulation with one or more  $\alpha_2$ -selective adrenergic agonists, carbonic anhydrase inhibitors or prostaglandin analogs. Examples of  $\alpha_2$ -selective adrenergic agonists include clonidine, apraclonidine, guanfacine, guanabenz and methyldopa, which are administered in effective amounts as is known in the art. Examples of carbonic anhydrase inhibitors include acetazolamide, dichlorphenamide and methazolamide, which are administered in effective amounts as is known in the art. Examples of prostaglandin analogs include PGE<sub>2</sub> and PGF<sub>2 $\alpha$</sub>  analogs, which are administered in effective amounts as is known in the art, including effective amounts administered by topical application to the eye. Thus, the invention further provides pharmaceutical compositions comprising an agent of the invention in combination with an effective amount of an  $\alpha_2$ -selective adrenergic agonist, carbonic anhydrase inhibitor, prostaglandin analog, or combination thereof.

Pharmaceutical compositions can be prepared to allow a therapeutically effective quantity of the compound of the present invention, and can include a pharmaceutically acceptable carrier, selected from known materials utilized for this purpose. *See, e.g.,* Remington, *The Science and Practice of Pharmacy*, 1995; *Handbook of Pharmaceutical Excipients*, 3<sup>rd</sup> Edition, 1999. Such compositions can be prepared in a variety of forms, depending on the method of administration.

In addition to the subject compound, the compositions of this invention can contain a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier", as used herein, means one or more compatible solid or liquid filler diluents or encapsulating substances that are suitable for administration to an animal, including a mammal or human. The term "compatible", as used herein, means that the components of the composition are capable of being commingled with the subject compound, and with each other, such that there is no interaction that would substantially reduce the

pharmaceutical efficacy of the composition under ordinary use. Preferably when liquid dose forms are used, the compounds of the invention are soluble in the components of the composition. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as corn starch and-potato starch; cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, and methyl cellulose; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid and magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerine, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the Tween<sup>TM</sup> brand emulsifiers; wetting agents, such sodium lauryl sulfate; coloring agents; flavoring agents; tableting agents, stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions. The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the subject compound is basically determined by the way the compound is to be administered. If the subject compound is to be injected, the preferred pharmaceutically-acceptable carrier is sterile, physiological saline, with a blood-compatible suspending agent, the pH of which has been adjusted to about 7.4.

If the preferred mode of administering the subject compound is perorally, the preferred unit dosage form is therefore tablets, capsules, lozenges, chewable tablets, and the like. Such unit dosage forms comprise a safe and effective amount of the subject compound, which is preferably from about 0.7 or 3.5 mg to about 280 mg/ 70 kg, more preferably from about 0.5 or 10 mg to about 210 mg/ 70 kg. The pharmaceutically-acceptable carrier suitable for the preparation of unit dosage forms for peroral administration are well-known in the art. Tablets typically comprise conventional pharmaceutically-compatible adjuvants as inert diluents, such as calcium carbonate, sodium carbonate, mannitol, lactose and cellulose; binders such as starch, gelatin and sucrose; disintegrants such as starch, alginic acid and croscarmellose; lubricants such as magnesium stearate, stearic acid and talc. Glidants such as silicon dioxide can be used to improve flow characteristics of the powder-mixture. Coloring agents, such as the FD&C

dyes, can be added for appearance. Sweeteners and flavoring agents, such as aspartame, saccharin, menthol, peppermint, and fruit flavors, are useful adjuvants for chewable tablets. Capsules typically comprise one or more solid diluents disclosed above. The selection of carrier components depends on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of this invention, and can be readily made by a person skilled in the art.

Peroral compositions also include liquid solutions, emulsions, suspensions, and the like. The pharmaceutically-acceptable carriers suitable for preparation of such compositions are well known in the art. Such liquid oral compositions preferably comprise from about 0.012% to about 0.933% of the subject compound, more preferably from about 0.033% to about 0.7%. Typical components of carriers for syrups, elixirs, emulsions and suspensions include ethanol, glycerol, propylene glycol, polyethylene glycol, liquid sucrose, sorbitol and water. For a suspension, typical suspending agents include methyl cellulose, sodium carboxymethyl cellulose, cellulose (e.g. Avicel™, RC-591), tragacanth and sodium alginate; typical wetting agents include lecithin and polyethylene oxide sorbitan (e.g. polysorbate 80). Typical preservatives include methyl paraben and sodium benzoate. Peroral liquid compositions may also contain one or more components such as sweeteners, flavoring agents and colorants disclosed above.

Other compositions useful for attaining systemic delivery of the subject compounds include sublingual and buccal dosage forms. Such compositions typically comprise one or more of soluble filler substances such as sucrose, sorbitol and mannitol; and binders such as acacia, microcrystalline cellulose, carboxymethyl cellulose and hydroxypropyl methyl cellulose. Glidants, lubricants, sweeteners, colorants, antioxidants and flavoring agents disclosed above may also be included.

Compositions can also be used to deliver the compound to the site where activity is desired; such as eye drops, gels and creams for ocular disorders.

Compositions of this invention include solutions or emulsions, preferably aqueous solutions or emulsions comprising a safe and effective amount of a subject compound intended for topical intranasal administration. Such compositions preferably comprise from about 0.01% to about 10.0% w/v of a subject compound, more preferably from about 0.1% to about 2.0%. Similar compositions are preferred for systemic delivery of subject compounds by the intranasal route. Compositions intended to deliver the compound systemically by intranasal dosing preferably comprise similar amounts of

a subject compound as are determined to be safe and effective by peroral or parenteral administration. Such compositions used for intranasal dosing also typically include safe and effective amounts of preservatives, such as benzalkonium chloride and thimerosal and the like; chelating agents, such as edetate sodium and others; buffers such as phosphate, citrate and acetate; tonicity agents such as sodium chloride, potassium chloride, glycerin, mannitol and others; antioxidants such as ascorbic acid, acetylcysteine, sodium metabisulfite and others; aromatic agents; viscosity adjustors, such as polymers, including cellulose and derivatives thereof; and polyvinyl alcohol and acids and bases to adjust the pH of these aqueous compositions as needed. The compositions may also comprise local anesthetics or other actives. These compositions can be used as sprays, mists, drops, and the like.

Other preferred compositions of this invention include aqueous solutions, suspensions, and dry powders comprising a safe and effective amount of a subject compound intended for atomization and inhalation administration. Such compositions are typically contained in a container with attached atomizing means. Such compositions also typically include propellants such as chlorofluorocarbons 12/11 and 12/114, and more environmentally friendly fluorocarbons, or other nontoxic volatiles; solvents such as water, glycerol and ethanol, these include cosolvents as needed to solvate or suspend the active; stabilizers such as ascorbic acid, sodium metabisulfite; preservatives such as cetylpyridinium chloride and benzalkonium chloride; tonicity adjustors such as sodium chloride; buffers; and flavoring agents such as sodium saccharin. Such compositions are useful for treating respiratory disorders, such as asthma and the like.

Other preferred compositions of this invention include aqueous solutions comprising a safe and effective amount of a subject compound intended for topical intraocular administration. Such compositions preferably comprise from about 0.01% to about 0.8% w/v of a subject compound, more preferably from about 0.05% to about 0.3%. Such compositions also typically include one or more of preservatives, such as benzalkonium chloride or thimerosal; vehicles, such as poloxamers, modified celluloses, povidone and purified water; tonicity adjustors, such as sodium chloride, mannitol and glycerin; buffers such as acetate, citrate, phosphate and borate; antioxidants such as sodium metabisulfite, butylated hydroxy toluene and acetyl cysteine; acids and bases can be used to adjust the pH of these formulations as needed.

Other preferred compositions of this invention useful for peroral administration include solids, such as tablets and capsules, and liquids, such as solutions, suspensions and emulsions (preferably in soft gelatin capsules), comprising a safe and effective amount of a subject compound. Such compositions can be coated by conventional methods, typically with pH or time-dependent coatings, such that the subject compound is released in the gastrointestinal tract at various times to extend the desired action. Such dosage forms typically include, but are not limited to, one or more of cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropyl methyl cellulose phthalate, ethyl cellulose, Eudragit<sup>TM</sup> coatings, waxes and shellac.

The compounds of the invention are administered by ocular, oral, parenteral, including, for example, using formulations suitable as eye drops. For ocular administration, ointments or droppable liquids may be delivered by ocular delivery systems known to the art such as applicators or eye droppers. Such compositions can include mucomimetics such as hyaluronic acid, chondroitin sulfate, hydroxypropyl methylcellulose or polyvinyl alcohol, preservatives such as sorbic acid, EDTA or benzylchromium chloride, and the usual quantities of diluents and/or carriers. See, Remington's Pharmaceutical Sciences, 16th Ed., Mack Publishing, Easton, PA, 1980, as well as later editions, for information on pharmaceutical compounding.

Numerous additional administration vehicles will be apparent to those of ordinary skill in the art, including without limitation slow release formulations, liposomal formulations and polymeric matrices.

In another preferred embodiment, the pharmaceutically effective amount is approximately 0.1 or 0.5 to 4 mg/kg body weight daily. Still more preferably, the pharmaceutically effective amount is approximately 1 mg/kg body weight daily. In a preferred embodiment, the amount is administered in once daily doses, each dose being approximately 1 mg/kg body weight.

Compounds of the invention can be used in conjunction with monitoring the improvement (decrease) in the intraocular pressure in a mammal using standard methodology.

The methods of the inventions can be assessed in animal models for ophthalmologic function. For example, improvements in fluid outflow facility can be studied in Rhesus monkeys treated with the compounds and methods of the invention. Aged Rhesus monkeys receive a single transcorneal injection of a test compound

(compound of the invention) at a concentration of about 1 mM in the anterior chamber of one eye, and Barany's solution, as a control, in the adjacent eye. Needle outflow facility is measured under baseline and pilocarpine-stimulated conditions at time points (for example, 3, 8, 12 and 24 weeks), after the administration of the test compound.

- 5 Increases in outflow facility in the drug treated vs. the control eye under baseline and cholinergic-stimulated (e.g. pilocarpine) conditions at the various time points are compared. As the enhancement of outflow facility can be influenced by the route of administration of the cholinergic agent, various routes of administration of the cholinergic agent can be used in the experiments. For instance, an intravenous  
10 administration versus a direct administration of pilocarpine can be compared. The above experiment demonstrates one method of measuring the improvement in ophthalmologic function. Such improvement has been illustrated with 4,5-dimethyl-3-(2-oxoethyl-phenethyl)thiazolium chloride, a compound believed to act by the same mechanism as those described here. See, U.S. application for "Methods for Treating Glaucoma I,"  
15 concurrently filed herewith.

In addition to measuring increased fluid outflow facility using the methods of the invention, improvements in pilocarpine-stimulated accommodation (i.e. the process of effecting refractive changes in the shape of the lens) can also be assessed in animal studies. As in the regulation of outflow facility, cholinergic input stimulates the  
20 movement of the ciliary muscle to control the shape of the lens, and allows accommodation in conditions of low illumination. Accommodation is impaired in a vast majority of individuals and begins to become noticeable to the individual around the age of 40 years. Interestingly, changes in accommodative response occur much earlier in life, around 18 years of age, and progresses until vision is noticeably impaired.

- 25 Physiological studies on accommodation are conducted following intraocular injection of a test compound and the results are compared relative to the results of control (untreated) animals. In the experiment, primates (for example, Rhesus monkeys) are treated twice a day for four days with 2  $\mu$ g of prostaglandin F<sub>2</sub> $\alpha$  (PGF<sub>2</sub> $\alpha$ ). On days 5-8 both eyes are treated first with 2  $\mu$ g of PGF<sub>2</sub> $\alpha$  followed 2 hours later with an  
30 intraocular injection of 10  $\mu$ L of the test compound of a final concentration of 1 mM. No injection is made to the control eye. 24 hours after the last injection of the test compound, a course of therapy consisting of once a day dosing for a total of 4 days

accommodative responses to i.m. pilocarpine administration is performed following phenylephrine refraction. Improvement in accommodation has been illustrated with 4,5-dimethyl-3-(2-oxoethyl-phenethyl)thiazolium chloride, a compound believed to act by the same mechanism as those described here. See, U.S. application for "Methods for

5 Treating Glaucoma I," concurrently filed herewith.

Compounds of the invention can be tested to determine corneal penetration to the anterior chamber of the eye following topical administration of eye drops. For example, a test compound is assayed *in vitro* through an intact rabbit cornea for transcorneal penetration in a standard diffusion chamber apparatus. Corneas are mounted in a

10 chamber at 37 °C with the epithelial side exposed to the test compound in Barany's solution. 1.0 mL samples are taken from the endothelial side 1 hour after addition of the test compound at a final concentration of 1 mM to the epithelial chamber. The volume of the chamber is replaced with phosphate buffered saline. The amount of test compound can be measured using any means that can be used to separate the compound

15 and measure its concentration. For example, an HPLC with an attached UV detector can be used to determine the concentration of the test compound that has penetrated the cornea. Penetration values are also determined at later time points, for example, at 5 hours.

Assessment of corneal penetration of compounds of the invention can be

20 determined *in vivo*, for example, in Cynomolgus monkeys. During these studies, the penetration of a test compound is evaluated using an eye-cup which holds a solution of 10 mM of the test compound in Barany's solution for 5 hours. At the end of the experiment the eye cup is removed, the eye is repeatedly flooded with Barany's solution and a sample of intraocular fluid is removed from the anterior chamber with a needle

25 inserted through the cornea. The quantity of the test compound in the intraocular fluid is determined using, for example, HPLC methods.

The activity of the compounds of the invention in breaking, reversing or inhibiting the formation of AGE's or AGE-mediated cross-links can be assayed by any of the methods described in US Patent 5,853,703.

30 The following examples further illustrate the present invention, but of course, should not be construed as in any way limiting its scope.



**EXAMPLE 1.** 2,6-diamino-benzothiazole dihydrochloride: 4 g of 2-amino-6-nitrobenzothiazole (Aldrich) was suspended in 130 ml MeOH, and 0.4 g 10% Pd/c (Aldrich) added. The suspension was hydrogenated at room temperature under 60 psi H<sub>2</sub> for 6.5 h. The reaction mixture was filtered, and the particulate washed with MeOH.

- 5 The filtrate was concentrated under reduced pressure, and crystals formed from the concentrate were collected to yield 2.67 g. mp 196-198°C, yield 81.6%. 0.91 g of this product was dissolved in 22 ml MeOH, and the pH adjusted with HCl to 4 to produce 1.2 g of crystals of 2,6-diamino-benzothiazole dihydrochloride. mp 318-320°C, 92.3% yield. Anal. calc. for C<sub>7</sub>H<sub>9</sub>N<sub>3</sub>SCl<sub>2</sub>, C 35.30%, H 3.80%, N 17.64%. Found, C 34.91%, H 3.67%, N 17.71%.
- 10

**EXAMPLE 2.** 2-(3,5-Dimethylphenoxy)-N-thiazol-2-yl)acetamide: First Route:

3,5-Dimethylphenol is reacted with bromoacetic acid at 110°C for four hours, with the reaction mixture stirred overnight without added heat. The resulting (3,5-dimethylphenoxy)acetic acid is dissolved in methylene chloride and coupled to 2-aminothiazole in an overnight, room temperature reaction conducted in the presence of base (N-methylmorpholine) and dehydration mediators 1-hydroxybenzotriazole and 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride.

- Second Route: 3,5-Dimethylphenol is reacted for 4.5 h with bromoacetic acid in THF under nitrogen and in the presence of sodium hydride. The resulting (3,5-dimethylphenoxy)acetic acid is reacted overnight with thionyl chloride, with heat. The resulting (3,5-dimethylphenoxy)acetyl chloride is reacted overnight with 2-aminothiazole in the presence of triethylamine, with cooling to 0°C.
- 20

- Third Route: 2-Aminothiazole (20g, 199.7 mmol) was suspended in methylene chloride (200 ml), in the presence of pyridine (20 ml, ~250 mmol), and the mixture cooled to 0°C. Bromoacetyl bromide (18.1 ml, 207.6 mmol) was dissolved in 400 ml methylene chloride, and this solution added to the suspended 2-aminothiazole dropwise. The resulting reaction mixture was stirred at room temperature overnight. The crude product was washed with water (200 ml, 1X), then sodium bicarbonate solution (200 ml, 2X), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The product
- 30
- 2-bromoacetamidothiazole was crystallized from MeOH. Yield, 4g. mp 148°C.

A solution of 3,5-dimethylphenol (2.5g, 13.9 mmol) in dry DMF (20 ml) was placed under a dry nitrogen atmosphere. Sodium hydride (0.7 g, 27.8 mmol; a 60% dispersion in mineral oil) was added in portions, and the mixture stirred for 1 h. A solution of 2-bromoacetamidothiazole (3.0 g, 13.9 mmol) in dry DMF (10 ml) was added to the mixture dropwise. The reaction was heated to 90°C for 5 h, then maintained overnight without external heat. The reaction mixture was poured into ice water, and the resulting material extracted with methylene chloride (50 ml X 3). The organic layer was washed with water (100 ml X 5), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The residue from evaporation was purified by silica gel chromatography developed with pet. ether: ether (1:1 v/v). The product N-(Thiazolyl)-2-(3,5-dimethylphenoxy)-acetamide was crystallized from acetonitrile and methyl tert-butyl ether. Yield, 1.04 g. mp 124-125 °C.

**EXAMPLE 3.** 2-Furyl-N-[4-(6-methyl-benzothiazol-2-yl)phenyl]carboxamide: First

Route: 2-Furoic acid (1.85 gms, 16.5 mmole) was dissolved in anhydrous methylene chloride (30 mls), to which solution was added a suspension of 2-(4-amino-phenyl)-6-methyl benzothiazole (4.76 g., 16.5 mmole) and N-methyl morpholine (2.0 g., 16.5 mmole) in methylene chloride (30 mls, at room temperature). Then, 1-hydroxy-benzotriazole hydrate (2.67 g., 16.5 mmole) and 1-(3-dimethyl amino propyl)-3-ethyl carbodiimide hydrochloride (4.75 g., 16.5 mmole) were added at room temperature. More methylene chloride (20 ml.) was added with stirring at room temperature, and the reaction maintained overnight. The initial clear reaction solution changed to a turbid solution. More methylene chloride (10 ml.) was added to the product mixture, which was then extracted with 1N HCl to separate a solid. The solid was filtered and washed with water. The product solid was crystallized from large amount of MeOH to yield 2.19 gm. (33.1%). mp. 238-240°C. <sup>1</sup>H and <sup>13</sup>C NMR were consistent with the expected product. TLC showed one spot (5% MeOH-CH<sub>2</sub>Cl<sub>2</sub> as developing solvent on silica gel plate).

Route 2: 2-(4-aminophenyl)-6-methyl benzothiazole (2.0 gm, 8.3 mmole) and 2-Furoyl chloride (1.086 gm., 8.32 mmole) were suspended in methylene chloride (30 ml, anhydrous). triethylamine (1.24 gm., 12.25 mmole) was added to the reaction mixture with stirring at room temperature for 2 days. (pH 7.0-7.2). Methylene chloride (50 ml)

- was added to the reaction mixture, and the reaction mixture extracted with 1 N HCl (50 ml) to separate a solid. The solid was filtered and washed with water to yield 1.3 gm. (46%) of the desired compound. The product was crystallized from MeOH to obtain 0.99 gm. mp. 238-240°C. <sup>1</sup>H and <sup>13</sup>C NMR were consistent with the expected product.
- 5 TLC showed one spot (5% MeOH-CH<sub>2</sub>Cl<sub>2</sub> as developing solvent on silica gel plate).

#### EXAMPLE 4. Cross-Linking Inhibition Assay

- The following method was used to evaluate the ability of the compounds to inhibit the cross-linking of glycated bovine serum albumin (AGE-BSA) to rat tail tendon collagen-coated 96-well plates.
- 10 AGE-BSA was prepared by incubating BSA at a concentration of 200 mg per ml with 200 mM glucose in 0.4M sodium phosphate buffer, pH 7.4 at 37°C for 12 weeks. The glycated BSA was then extensively dialyzed against phosphate buffer solution (PBS) for 48 hours with additional 5 times buffer exchanges. The rat tail tendon
- 15 collagen coated plate was blocked first with 300 microliters of Superbloc blocking buffer (Pierce Chemical, Rockford, IL) for one hour. The blocking solution was removed from the wells by washing the plate twice with phosphate buffered saline (PBS)-Tween 20 solution (0.05% Tween 20) using a NUNC-multiprobe (Nalge Nunc, Rochester, NY) or Dynatech ELISA-plate (Dynatech, Alexandria, VA) washer. Cross-linking of AGE-BSA
- 20 (1 to 10 microgram per well depending on the batch of AGE-BSA) to rat tail tendon collagen coated plate was performed with and without the testing compound dissolved in PBS buffer at pH 7.4 at one or more desired concentrations by the addition of 50 microliters each of the AGE-BSA diluted in PBS or in the solution of test compound at 37°C for 4 hours. Unbrowned BSA in PBS buffer with or without testing compound
- 25 were added to the separate wells as the blanks. The un-cross-linked AGE-BSA was then removed by washing the wells three times with PBS-Tween buffer. The amount of AGE-BSA crosslinked to the tail tendon collagen-coated plate was then quantitated using a polyclonal antibody raised against AGE-RNase. After a one-hour incubation period, AGE antibody was removed by washing 4 times with PBS-Tween.
- 30 The bound AGE antibody was then detected with the addition of horseradish peroxidase-conjugated secondary antibody—goat anti-rabbit immunoglobulin and incubation for 30 minutes. The substrate of 2,2-azino-di(3-ethylbenzthiazoline sulfonic

acid) (ABTS chromogen) (Zymed Laboratories, Inc., South San Francisco, CA) was added. The reaction was allowed for an additional 15 minutes and the absorbance was read at 410 nm in a Dynatech plate reader.

## 5 EXAMPLE 5 Cross-Link Breaking Assay

To ascertain the ability of the compounds of the instant invention to break or reverse already formed advanced glycosylation endproducts, a sandwich enzyme immunoassay was applied. Generally, the assay utilizes collagen-coated 96 well microtiter plates that are obtained commercially. AGE-modified protein (AGE-BSA) is incubated on the collagen-coated wells for four hours, is washed off the wells with PBS-Tween and solutions of the test compounds are added. Following an incubation period of 16 hours (37°C) cross-link-breaking is detected using an antibody raised against AGE-ribonuclease or with an antibody against BSA.

### Preparation of solutions and buffers

- 15 Bovine Serum Albumin (Type V) (BSA) (from Calbiochem) solution was prepared as follows: 400 mg of Type V BSA (bovine serum albumin) was added for each ml of 0.4 M sodium phosphate buffer, pH 7.4. A 400 mM glucose solution was prepared by dissolving 7.2 grams of dextrose in 100 ml of 0.4 M sodium phosphate buffer, pH 7.4. The BSA and glucose solutions were mixed 1:1 and incubated at 37°C for 12 weeks.
- 20 The pH of the incubation mixture was monitored weekly and adjusted to pH 7.4 if necessary. After 12 weeks, the AGE-BSA solution was dialyzed against PBS for 48 hours with four buffer changes, each at a 1:500 ratio of solution to dialysis buffer. Protein concentration was determined by the micro-Lowry method. The AGE-BSA stock solution was aliquoted and stored at -20°C.
- 25 Test compounds were dissolved in PBS and the pH was adjusted to pH 7.4, if necessary. AGE-BSA stock solution was diluted in PBS to measure maximum crosslinking and in the inhibitor solution for testing inhibitory activity of compounds. The concentration of AGE-BSA necessary to achieve the optimum sensitivity was determined by initial titration of each lot of AGE-BSA.
- 30 Substrates for detection of secondary antibody binding were prepared by diluting the HRP substrate buffer (Zymed) 1:10 in distilled water and mixing with ABTS chromogen (Zymed) 1:50 just prior to use.

Assay Procedures

Biocoat plates were blocked with 300 microliters of Superbloc (Pierce Chemical). Plates were blocked for one hour at room temperature and were washed with PBS-Tween (0.05% v/v) three times with the Dynatech platewasher before addition of test reagents.

- 5 The first three wells of the Biocoat plate were used for the reagent blank. Fifty microliters of solutions AGE-BSA were added to test wells in triplicate and only PBS in blank wells. The plate was incubated at 37°C for four hours and washed with PBS-Tween three times. Fifty microliters of PBS was added to the control wells and 50 microliters of the test prospective agent was added to the test wells and blank. The plate  
10 was incubated overnight (approximately 16 hours) with prospective agent, followed by washing in PBS before addition of primary antibody.

(Prior to use, each lot of primary antibody, either anti-BSA or anti-RNase, was tested for optimum binding capacity in this assay by preparing serial dilutions (1:500 to 1:2000) and plating 50 microliters of each dilution in the wells of Biocoat plates.

- 15 Optimum primary antibody was determined from saturation kinetics.) Fifty microliters of primary antibody of appropriate dilution, was added and incubated for one hour at room temperature. The plate was then washed with PBS-Tween.

- Plates were incubated with the secondary antibody, HRP-(Goat-anti-rabbit), which was diluted 1:4000 in PBS and used as the final secondary antibody. The  
20 incubation was performed at room temperature for thirty minutes.

Detection of maximum crosslinking and breaking of AGE crosslinking was performed as follows. HRP substrate (100 microliter) was added to each well of the plate and was incubated at 37°C for fifteen minutes. Readings were taken in the Dynatech ELISA-plate reader.

25

Definitions

**Heterocycle.** Except where heteroaryl is separately recited for the same substituent, the term "heterocycle" includes heteroaryl.

- All publications and references, including but not limited to patents and patent  
30 applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent

application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

While this invention has been described with an emphasis upon preferred embodiments, it will be obvious to those of ordinary skill in the art that variations in the preferred devices and methods may be used and that it is intended that the invention may be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the claims that follow.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
22